

International Journal of Pure and Applied Mathematics - IJPAM

ISSN 1311-8080

[links](#)

[contact](#)

Indexing and Abstracting

"International Electronic Journal of Pure and Applied Mathematics" is indexed/abstracted in:

- [ELSEVIER](#);
- [Index Copernicus](#);
- [Hub SciVerse](#);
- [Google Scholar](#);
- [SCImago Journal & Country Rank](#);
- [SCIRUS](#);
- [SCOPUS](#);
- [Zentralblatt MATH](#).

We would like to thank these institutions for cooperation and support.

Search

Loading

IJPAM

[Academic Publications, Ltd.](#)

[Home](#)


[Contents](#)

[Editorial Board](#)

[How to?](#)

[Instruction for Authors](#)

[Subscriptions](#)



International Journal of Pure and Applied Mathematics

ISSN 1311-8080 (PRINTED VERSION)
ISSN 1314-3395 (ON-LINE VERSION)

[links](#) [contact](#)

Contents IJPAM: Volume 84, No. 4 (2013)

M.M. Khader, N.H. Sweilam, T.A. Assiri

ON THE NUMERICAL SOLUTION FOR THE FRACTIONAL WAVE EQUATION USING LEGENDRE PSEUDOSPECTRAL METHOD ... **307**

[| more](#) | [download this paper \(free\)](#) |

K. Aruna, A.S.V. Ravi Kanth

A NOVEL APPROACH FOR A CLASS OF HIGHER ORDER NONLINEAR SINGULAR BOUNDARY VALUE PROBLEMS ... **321**

[| more](#) | [download this paper \(free\)](#) |

R. Behl, V. Kanwar, K.K. Sharma

MODIFIED OPTIMAL FAMILIES OF FOURTH-ORDER JARRATT'S METHOD ... **331**

[| more](#) | [download this paper \(free\)](#) |

H. Kim

THE SOLUTION OF EULER-CAUCHY EQUATION EXPRESSED BY DIFFERENTIAL OPERATOR USING LAPLACE TRANSFORM ... **345**

[| more](#) | [download this paper \(free\)](#) |

A.K. Dubey, S.K. Tiwari, R.P. Dubey

CONE METRIC SPACES AND FIXED POINT THEOREMS OF GENERALIZED T-CONTRACTIVE MAPPINGS ... **353**

[| more](#) | [download this paper \(free\)](#) |

K. Teerapabolarn

AN IMPROVED BOUND ON GEOMETRIC APPROXIMATION BY w -FUNCTIONS ... **365**

[| more](#) | [download this paper \(free\)](#) |

W.A. Khan, A. Taouti

ON P-KRULL MONOIDS ... **371**

[| more](#) | [download this paper \(free\)](#) |

V. Arasu, J.B. Stalin

GENERALIZED HYERS-ULAM-RASSIAS THEOREM OF QUADRATIC FUNCTIONAL EQUATION IN MENGER PROBABLISTIC NORMED SPACES ... **377**

[| more](#) | [download this paper \(free\)](#) |

H. Arsham

MODELING TARIFF TO ANALYZE PROTECTIONISM ... **389**

[| more](#) | [download this paper \(free\)](#) |

V. Oussa

SHANNON-LIKE PARSEVAL FRAME WAVELETS ON SOME TWO STEP NILPOTENT LIE GROUPS ... **397**

[| more](#) | [download this paper \(free\)](#) |

K. Teerapabolarn

A NEW BOUND ON POISSON APPROXIMATION FOR INDEPENDENT GEOMETRIC VARIABLES ... **419**

[| more](#) | [download this paper \(free\)](#) |

A. Selvam, P.S. Kala, N. Marikkannan

ON APPLICATIONS OF DIFFERENTIAL SUBORDINATION ... **423**

[| more](#) | [download this paper \(free\)](#) |

B. Turmetov, K. Shinaliyev

ON A METHOD FOR CONSTRUCTING SOLUTIONS OF DIFFERENTIAL EQUATIONS OF FRACTIONAL ORDER WITH A HADAMARD TYPE OPERATOR ... **435**

[| more](#) | [download this paper \(free\)](#) |

Search

Google™ Custom Search ×
powered by Google™

IJPAM

[Home](#)

[Contents](#)

[Editorial Board](#)

[How to?](#)

[Indexing and Abstracting](#)

[Instruction for Authors](#)

[Subscriptions](#)



**ON A METHOD FOR CONSTRUCTING SOLUTIONS OF
DIFFERENTIAL EQUATIONS OF FRACTIONAL
ORDER WITH A HADAMARD TYPE OPERATOR**

Batirkhan Turmetov¹, Kanat Shinaliyev^{2 §}

Department of Mathematics

International Kazakh-Turkish

University named after A. Yasawi

Turkestan, 161200, REPUBLIC OF KAZAKHSTAN

Abstract: In this paper we investigate a method for constructing solutions of differential equations of fractional order with Hadamard type derivative. The basis of this method is the construction of systems concerning the operator of fractional differentiation with Hadamard type derivative. We define new classes of special functions generalizing the Mittag-Leffler functions, and related to the differential equations of fractional order. Properties of these functions and their applications to solutions of fractional order differential equations are established. Analytical solutions of differential equations of fractional order are found as a series. Moreover, we consider normed systems of non-homogeneous differential equations of fractional order.

AMS Subject Classification: 35J40, 35J05

Key Words: fractional derivative, Hadamard operator, normed system, operator method

Introduction

Let $0 < a < b < \infty$, $\alpha > 0$. For a given α the following expression is called an

Received: January 21, 2013

© 2013 Academic Publications, Ltd.
url: www.acadpubl.eu

[§]Correspondence author

integration operator of α order in the sense of Hadamard[1]:

$$I_a^\alpha(f)(t) = \frac{1}{\Gamma(\alpha)} \int_a^t \left(\ln \frac{t}{\tau}\right)^{\alpha-1} \frac{f(\tau)}{\tau} d\tau, \quad t > a.$$

Furthermore, we put $I_a^0(f)(t) \equiv f(t)$.

Let $\delta = t \frac{d}{dt}$, $\delta^k = \delta(\delta^{k-1})$, $k = 1, 2, \dots$

Then for $\alpha \in (m-1, m]$, $m = 1, 2, \dots$ the following expressions are called differentiation operators of α in the sense of, respectively, Hadamard and Hadamard-Caputo, see [1]:

$${}_H D_a^\alpha[f](t) = \delta^m (I_a^{m-\alpha} f)(t),$$

$${}_{HC} D_a^\alpha[f](t) = I_a^{m-\alpha} (\delta^m f)(t)$$

Let $0 \leq \beta \leq 1$, $m-1 < \alpha \leq m$, $m = 1, 2, \dots$. Then operator

$$D_a^{\alpha,\beta}(f)(t) = I^{\beta(m-\alpha)} \delta^m I^{(1-\beta)(m-\alpha)}(f)(t)$$

is called differentiation operator of α order and β type (see [2, 3]). It is clear that if $\beta = 0$ then the operator $D_a^{\alpha,\beta}$ coincides with ${}_H D_a^\alpha$, and when $\beta = 1$ with the operator ${}_{HC} D_a^\alpha$. Note that properties and applications of the operator $D_a^{\alpha,\beta}$ when $0 < \alpha \leq 1$, $0 \leq \beta \leq 1$ have been studied in [2].

In this paper we study a method for constructing solutions of differential equations of the following form:

$$\left(D_a^{\alpha,\beta} - \lambda\right)^n y(t) = h(t). \quad (1)$$

where $\lambda \in R$, $n = 1, 2, \dots$. This method was studied for the differential equations of integer order in [4], and for the equations of fractional order with Riemann-Liouville type operator in [5, 6]. The basis of this method is the construction of normed systems for the operators $D_a^{\alpha,\beta}$ and $D_a^{\alpha,\beta} - \lambda$.

Consider definition of the normed systems [4]. Let L_1 and L_2 be linear operators given in a linear space of functions X . An infinitely system of functions $f_k(x) \in X$, $k = 1, 2, \dots$, where any finite subsystem is linear independent, is called f -normed concerning (L_1, L_2) with base $f_0(x)$, if everywhere in this domain the following equality holds:

$$L_1 f_0(x) = f(x), L_1 f_k(x) = L_2 f_{k-1}(x), k = 1, 2, \dots$$

A particular case of this definition is the case when L_2 is an identity operator. In this case, instead of normalizability on the pair of operators (L_1, L_2)

we will talk about normalizability on one operator L_1 . If $f(x) = 0$, then the system $f_k(x) \in X, k = 1, 2, \dots$ is called 0-normed or just normed [4].

An important property of the normed system is the following: if $\{f_k(x)\}$ is a system of functions f -normed concerning (L_1, L_2) in the domain Ω such that the series $y(x) = \sum_{k=0}^{\infty} f_k(x)$ converges, and admits termwise application of operators L_1 and L_2 , then the function $y(x)$ in the domain Ω is a solution of the equation:

$$(L_1 - L_2) y(x) = f(x).$$

It is easy to check that if $\{f_k(x)\}$ is a system of functions f -normed concerning L_1 in the domain Ω , and the following equalities hold:

$$L_1 L_2 f_k = L_2 L_1 f_k, k = 1, 2, \dots$$

Then the system of functions $\{L_2^k f_k(x)\}$ is f -normed concerning (L_1, L_2) in the domain Ω . Therefore, to use the basic property of the normed system of functions during the construction of solutions of equation $(L_1 - L_2) y(x) = f(x)$ it is necessary to find f -normed in Ω system of functions concerning the operator L_1 .

1. Construction of Normed System for Homogenous Equation

Let $0 \leq \beta \leq 1, m - 1 < \alpha \leq m, m = 1, 2, \dots, \gamma = (1 - \beta)(m - \alpha)$. Introduce the coefficients:

$$C(\alpha, \beta, s, 0) = 1, C(\alpha, \beta, s, i) = \frac{\Gamma(i\alpha + s - \delta + 1)}{\Gamma(s - \delta + 1)}, i \geq 1,$$

where $s = 0, 1, \dots, m - 1$. Let us consider some properties of the operators I_a^α and $D_a^{\alpha, \beta}$.

Lemma 1. *Let $s = 0, 1, \dots, m - 1, m \geq 1$, then*

$$\delta^m \left(\ln \frac{t}{a} \right)^s = 0. \tag{2}$$

Proof. It is obvious that $\delta \left(\ln \frac{t}{a} \right)^0 = 0$ and $\delta \left(\ln \frac{t}{a} \right) = 1$. Then for all $s \geq 1$:

$$\delta \left(\ln \frac{t}{a} \right)^s = s \cdot \left(\ln \frac{t}{a} \right)^{s-1}.$$

Further, by induction it easy follows that

$$\delta^k \left(\ln \frac{t}{a} \right)^s = s \cdot (s-1) \cdot \dots \cdot (s-k+1) \left(\ln \frac{t}{a} \right)^{s-k}, \quad s \geq k, \quad (3)$$

and

$$\delta^k \left(\ln \frac{t}{a} \right)^s = 0, \text{ if } k > s.$$

Consequently, for all $s = 0, 1, \dots, m-1$ equality (2) holds. Lemma is proved.

Lemma 2. Let $\mu > -1$ and $f(t) = \left(\ln \frac{t}{a} \right)^\mu$. Then

$$I_a^\alpha(f)(t) = \frac{\Gamma(\mu+1)}{\Gamma(\alpha+\mu+1)} \left(\ln \frac{t}{a} \right)^{\alpha+\mu}. \quad (4)$$

Proof. By definition of the operator I_a^α we have

$$\begin{aligned} I_a^\alpha(f)(t) &= \frac{1}{\Gamma(\alpha)} \int_a^t \left(\ln \frac{t}{\tau} \right)^{\alpha-1} \left(\ln \frac{\tau}{a} \right)^\mu \frac{d\tau}{\tau} = \\ &= \frac{1}{\Gamma(\alpha)} \int_a^t \left(\ln \frac{t}{a} - \ln \frac{\tau}{a} \right)^{\alpha-1} \left(\ln \frac{\tau}{a} \right)^\mu d \ln \frac{\tau}{a} = \\ &= \frac{1}{\Gamma(\alpha)} \int_0^1 (1-\xi)^{\alpha-1} \xi^\mu d\xi \left(\ln \frac{t}{a} \right)^{\alpha+\mu} = \frac{\Gamma(\mu+1)}{\Gamma(\alpha+\mu+1)} \left(\ln \frac{t}{a} \right)^{\alpha+\mu}. \end{aligned}$$

Lemma is proved.

Lemma 3. Let $\mu > m-1$ and $f(t) = \left(\ln \frac{t}{a} \right)^\mu$. Then

$$D_a^{\alpha,\beta}(f)(t) = \frac{\Gamma(\mu+1)}{\Gamma(\mu+1-\alpha)} \left(\ln \frac{t}{a} \right)^{\mu-\alpha}. \quad (5)$$

Proof. Let $\mu > m-1$. Due to the equality (4) for the functions $I_a^\gamma(f)(t)$ we get

$$I_a^\gamma(f)(t) = \frac{\Gamma(\mu+1)}{\Gamma(\mu+1+\gamma)} \left(\ln \frac{t}{a} \right)^{\mu+\gamma}.$$

Then by the equality (3) we have

$$\begin{aligned} \delta^m I_a^\gamma(f)(t) &= \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1 + \gamma)} \delta^m \left(\ln \frac{t}{a}\right)^{\mu + \gamma} = \\ &= \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1 + \gamma)} (\mu + \gamma) \dots (\mu + \gamma - m + 1) \left(\ln \frac{t}{a}\right)^{\mu + \gamma - m} = \\ &= \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1 + \gamma - m)} \left(\ln \frac{t}{a}\right)^{\mu + \gamma - m}. \end{aligned}$$

Further,

$$I_a^{\beta(m-\alpha)} \left(\ln \frac{t}{a}\right)^{\mu + \gamma - m} = \frac{\Gamma(\mu + \gamma + 1 - m)}{\Gamma(\mu + \gamma + 1 - m + \beta(m - \alpha))} \left(\ln \frac{t}{a}\right)^{\mu + \gamma - m + \beta(m - \alpha)}.$$

Since $\gamma + \beta(m - \alpha) = m - \alpha$, then

$$\begin{aligned} D_a^{\alpha, \beta}(f)(t) &= I_a^{\beta(m-\alpha)} \delta^m I_a^\gamma(f)(t) = \\ &= \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1 + \gamma - m)} \frac{\Gamma(\mu + 1 + \gamma - m)}{\Gamma(\mu + 1 - \alpha)} \left(\ln \frac{t}{a}\right)^{\mu - \alpha} = \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1 - \alpha)} \left(\ln \frac{t}{a}\right)^{\mu - \alpha} \end{aligned}$$

Lemma is proved.

Corollary 1. Let $\mu = k\alpha + s - \gamma$, $k = 1, 2, \dots$, $s = 0, 1, \dots, m - 1$ and $f(t) = \left(\ln \frac{t}{a}\right)^\mu$. Then

$$D_a^{\alpha, \beta}(f)(t) = \frac{\Gamma(k\alpha + s - \gamma + 1)}{\Gamma((k - 1)\alpha + s - \gamma + 1)} \left(\ln \frac{t}{a}\right)^{k\alpha + s - \gamma - \alpha}. \tag{6}$$

Lemma 4. Suppose that $f_i(t) = \frac{1}{C(\alpha, \beta, s, i)} \left(\ln \frac{t}{a}\right)^{\alpha i + s - \gamma}$, $i = 0, 1, \dots$. Then $f_i(t)$ for any $s = 0, 1, \dots, m - 1$ forms 0-normed system of the functions concerning the operator $D_a^{\alpha, \beta}$.

Proof. Let $i \geq 1$. Then, using (6) and definition of the coefficients $C(\alpha, \beta, s, i)$, we get

$$\begin{aligned} D_a^{\alpha, \beta} f_i(t) &= \frac{\Gamma(i\alpha + s - \gamma + 1)}{\Gamma((i - 1)\alpha + s - \gamma + 1)} \cdot \frac{\Gamma(s + 1 - \gamma)}{\Gamma(\alpha i + s - \gamma + 1)} \left(\ln \frac{t}{a}\right)^{\alpha i + s - \gamma - \alpha} = \\ &= \frac{\Gamma(s + 1 - \gamma)}{\Gamma((i - 1)\alpha + s - \gamma + 1)} \left(\ln \frac{t}{a}\right)^{(i-1)\alpha + s - \gamma} = \end{aligned}$$

$$= \frac{1}{C(\alpha, \beta, s, i - 1)} \left(\ln \frac{t}{a} \right)^{(i-1)\alpha+s-\gamma} = f_{i-1}(t).$$

If $i = 0$, then, by the equality (4), we have:

$$I_a^\gamma \left(\ln \frac{t}{a} \right)^{s-\gamma} = \frac{\Gamma(s - \gamma + 1)}{\Gamma(s - \gamma + 1 + \alpha)} \left(\ln \frac{t}{a} \right)^s.$$

Since

$$\delta^m \left(\ln \frac{t}{a} \right)^s = 0, \quad s = 0, 1, \dots, m - 1,$$

then

$$D_a^{\alpha, \beta} \left(\ln \frac{t}{a} \right)^{s-\gamma} = 0.$$

Lemma is proved.

Consider the function:

$$\Phi_{\alpha, \beta, s}^p(t) = \sum_{i=p}^{\infty} \lambda^{i-p} \binom{i}{p} \frac{t^{\alpha i + s - \delta}}{C(\alpha, \beta, s, i)},$$

where

$$p = 0, 1, \dots, \quad \binom{i}{p} = \frac{i!}{p!(i-p)!}.$$

Introduce the following spaces:

$$C_{\gamma, \ln}[a, b] = \left\{ y(t) : \left(\ln \frac{t}{a} \right)^\gamma \cdot y(t) \in C[a, b] \right\},$$

and

$$C_\gamma^\mu[a, b] = \left\{ y(t) \in C_{\gamma, \ln}[a, b] : D_a^{\alpha, \beta} y(t) \in C_{\mu, \ln}[a, b] \right\}, \mu > 0.$$

It is obvious that the spaces $C_{\gamma, \ln}[a, b]$ and $C_\gamma^\mu[a, b]$ are linear. The basic properties of the functions $\Phi_{\alpha, \beta, s}^p(\lambda, t)$ are presented in the following theorem:

Theorem 1. *Let $m - 1 < \alpha \leq m, 0 \leq \beta \leq 1, s = 0, 1, \dots, m - 1$. Then $\Phi_{\alpha, \beta, s}^p(\lambda, t) \in C_{\gamma, \ln}^\gamma[a, b]$, and for all $p = 0, 1, \dots$, forms 0-normed system concerning operator $D_a^{\alpha, \beta}$ i.e. the following equalities hold:*

$$\left(D_a^{\alpha, \beta} - \lambda \right) \Phi_{\alpha, \beta, s}^0(\lambda, t) = 0, \tag{7}$$

$$\left(D_a^{\alpha, \beta} - \lambda \right) \Phi_{\alpha, \beta, s}^p(\lambda, t) = \Phi_{\alpha, \beta, s}^{p-1}(\lambda, t), \quad p \geq 1. \tag{8}$$

Proof. Let $p = 0$. Then the function $\Phi_{\alpha,\beta,s}^0(\lambda, t)$ can be presented as a form:

$$\Phi_{\alpha,\beta,s}^0(\lambda, t) = \left(\ln \frac{t}{a}\right)^{-\gamma} \sum_{i=0}^{\infty} \frac{\lambda^i}{C(\alpha, \beta, s, i)} \left(\ln \frac{t}{a}\right)^{\alpha i+s} = \left(\ln \frac{t}{a}\right)^{-\gamma} y_1(t),$$

where

$$y_1(t) = \sum_{i=0}^{\infty} \frac{\lambda^i}{C(\alpha, \beta, s, i)} \left(\ln \frac{t}{a}\right)^{\alpha i+s}. \tag{9}$$

It is known that for a gamma function the following asymptotical estimate holds:

$$\Gamma(z + 1) = \sqrt{2\pi z} \left(\frac{z}{e}\right)^z \left\{1 + O\left(\frac{1}{z}\right)\right\}, \quad z \rightarrow \infty.$$

Then it is easy to show that the series (9) converges for $t \in [a, b]$, and the function $y_1(t) \in C[a, b]$. Consequently, $\Phi_{\alpha,\beta,s}^0(\lambda, t) \in C_\gamma[a, b]$. Using operator $D_a^{\alpha,\beta}$ to the function $\Phi_{\alpha,\beta,s}^0(\lambda, t)$, and taking account that the system $f_i(t) = \frac{1}{C(\alpha,\beta,s,i)} \left(\ln \frac{t}{a}\right)^{\alpha i+s}$ forms 0-normed system concerning operator $D_a^{\alpha,\beta}$, we receive

$$\begin{aligned} D_a^{\alpha,\beta} [\Phi_{\alpha,\beta,s}^0](\lambda, t) &= \sum_{i=1}^{\infty} \frac{\lambda^i}{C(\alpha, \beta, s, i-1)} \left(\ln \frac{t}{a}\right)^{(i-1)\alpha+s} = \\ &= \sum_{i=0}^{\infty} \frac{\lambda^{i+1}}{C(\alpha, \beta, s, i)} \left(\ln \frac{t}{a}\right)^{\alpha i+s} = -\lambda \Phi_{\alpha,\beta,s}^0(\lambda, t). \end{aligned}$$

Therefore, $D_a^{\alpha,\beta} [\Phi_{\alpha,\beta,s}^0](\lambda, t) \in C_\gamma^\gamma[a, b]$, and it satisfies equality (7). Now let $p \geq 1$. We use operator $D_a^{\alpha,\beta}$ to the function $\Phi_{\alpha,\beta,s}^p(\lambda, t)$, and similarly as in the case $p = 0$, we get

$$\begin{aligned} D_a^{\alpha,\beta} [\Phi_{\alpha,\beta,s}^p](\lambda, t) &= \sum_{i=p}^{\infty} \lambda^{i-p} \binom{i}{p} D_a^{\alpha,\beta} f_i(t) = \\ &= \sum_{i=p}^{\infty} \lambda^{i-p} \binom{i}{p} \frac{1}{C(\alpha, \beta, s, i)} \left(\ln \frac{t}{a}\right)^{\alpha(i-1)+s-\delta}. \end{aligned}$$

Further, in the last expression by changing the summation index and using the equality:

$$\binom{i+1}{p} - \binom{i}{p-1} = \binom{i}{p} \text{ and } \binom{p}{p} - \binom{p-1}{p-1} = 0,$$

we obtain

$$\begin{aligned}
 & D_a^{\alpha,\beta} \Phi_{\alpha,\beta,s}^p(\lambda, t) - \Phi_{\alpha,\beta,s}^{p-1}(\lambda, t) = \\
 &= \sum_{i=p-1}^{\infty} \frac{\lambda^{j-(p-1)}}{C(\alpha, s, j)} \left[\binom{i+1}{p} - \binom{i}{p-1} \right] \left(\ln \frac{t}{a} \right)^{\alpha j + s - \delta} = \\
 &= \sum_{i=p-1}^{\infty} \frac{\lambda^{j-(p-1)}}{C(\alpha, s, j)} \binom{i}{p} \left(\ln \frac{t}{a} \right)^{\alpha j + s - \delta} = \lambda \Phi_{\alpha,\beta,s}^p(\lambda, t)
 \end{aligned}$$

It follows that

$$(D_a^{\alpha,\beta} - \lambda) \Phi_{\alpha,\beta,s}^p(\lambda, t) = \Phi_{\alpha,\beta,s}^{p-1}(\lambda, t), \quad p \geq 1.$$

It is obvious that $\Phi_{\alpha,\beta,s}^p(\lambda, t) \in C[a, b]$ for all $p = 1, \dots, n-1$, and $\Phi_{\alpha,\beta,s}^0(\lambda, t) \in C_\gamma[a, b]$. Hence $D_a^{\alpha,\beta} \Phi_{\alpha,\beta,s}^p(\lambda, t) \in C_\gamma[a, b]$. Theorem is proved.

Corollary 2. *Suppose that conditions of theorem 1 hold for all $p = 0, 1, 2, \dots, n-1$ and $s = 0, 1, \dots, m-1$. Then the functions $\Phi_{\alpha,\beta,s}^p(\lambda, t)$ satisfy equality (1) when $h(t) = 0$.*

Let $p = 0$. Transform the function $\Phi_{\alpha,\beta,s}^0(\lambda, t)$ to the following form:

$$\begin{aligned}
 \Phi_{\alpha,\beta,s}^0(\lambda, t) &= \Gamma(s - \gamma + 1) \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + s - \gamma + 1)} \left(\ln \frac{t}{a} \right)^{\alpha i + s - \delta} = \\
 &= \Gamma(s - \gamma + 1) \left(\ln \frac{t}{a} \right)^{s - \delta} E_{\alpha, s - \gamma + 1} \left(\lambda \left(\ln \frac{t}{a} \right)^\alpha \right),
 \end{aligned}$$

where

$$E_{\alpha,\rho}(z) = \sum_{i=0}^{\infty} \frac{z^i}{\Gamma(i\alpha + \rho)}$$

is a Mittag-Leffler type function, see [1].

2. Construction of Normed System for Non-Homogenous Equation

Consider the function

$$h_0(t) = \int_a^t \left(\ln \frac{t}{\xi} \right)^{\alpha-1} E_{\alpha,\alpha} \left(\lambda \left(\ln \frac{t}{\xi} \right)^\alpha \right) \frac{h(\xi)}{\xi} d\xi. \tag{10}$$

In further we assume that $h(t)$ belongs to a class of functions for which all the transactions in question are valid. Let us show that

$$\left(D_a^{\alpha,\beta} - \lambda\right) h_0(t) = h(t). \tag{11}$$

Consider the case when $\alpha \in (0, 1]$. In this case by definition of the operator $D_a^{\alpha,\beta}$ we have

$$D_a^{\alpha,\beta} h_0(t) = \frac{1}{\Gamma(\gamma_1)} \cdot \int_a^t \left(\ln \frac{t}{\tau}\right)^{\gamma_1-1} \tau \frac{d\tau}{\tau} I_a^\gamma[h](\tau) \frac{d\tau}{\tau},$$

where $\gamma_1 = \beta(1 - \alpha)$.

Then

$$\begin{aligned} D_a^{\alpha,\beta} h_0(t) &= \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \frac{1}{\gamma_1} \left(\ln \frac{t}{\tau}\right)^{\gamma_1} \frac{d}{d\tau} I_a^\gamma[h](\tau) d\tau \right\} = \\ &= \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \frac{1}{\gamma_1} \left(\ln \frac{t}{\tau}\right)^{\gamma_1} I_a^\gamma[h](\tau) \Big|_{\tau=a}^{\tau=t} + \int_a^t \left(\ln \frac{t}{\tau}\right)^{\gamma-1} I_a^\gamma[h](\tau) d\tau \right\} = \\ &= \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ - \int_a^t \left(\ln \frac{t}{\tau}\right)^{\gamma_1-1} \frac{1}{\Gamma(\gamma)} \int_a^\tau \left(\ln \frac{\tau}{\xi}\right)^{\gamma-1} h_0(\xi) \frac{d\xi}{\xi} \frac{d\tau}{\tau} \right\} = \\ &= \frac{1}{\Gamma(\gamma_1)\Gamma(\gamma)} t \frac{d}{dt} \int_a^t h_0(\xi) \int_\xi^t \left(\ln \frac{t}{\tau}\right)^{\gamma_1-1} \left(\ln \frac{\tau}{\xi}\right)^{\gamma-1} \frac{d\tau}{\tau} \frac{d\xi}{\xi}. \end{aligned}$$

Let us study inner integral. We get

$$\begin{aligned} &\int_\xi^t \left(\ln \frac{t}{\xi} - \ln \frac{\tau}{\xi}\right)^{\gamma_1-\alpha} \left(\ln \frac{\tau}{\xi}\right)^{\gamma-1} d \ln \frac{\tau}{\xi} = \\ &= \int_0^1 (1-z)^{\gamma_1-1} z^{\gamma-1} dz \cdot \left(\ln \frac{t}{\xi}\right)^{\gamma+\gamma_1-1} = \frac{\Gamma(\gamma_1)\Gamma(\gamma)}{\Gamma(\gamma+\gamma_1)} \left(\ln \frac{t}{\xi}\right)^{\gamma+\gamma_1-1}. \end{aligned}$$

Further, by the presentation of the function $h_0(\xi)$ and taking account that $\gamma + \gamma_1 = 1 - \alpha$, we have

$$\begin{aligned} & \int_a^t h_0(\xi) \left(\ln \frac{t}{\xi}\right)^{\gamma+\gamma_1-1} \frac{d\xi}{\xi} = \\ &= \int_a^t \left(\ln \frac{t}{\xi}\right)^{-\alpha} \int_a^{\xi} \left(\ln \frac{\xi}{\tau}\right)^{\alpha-1} \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + \alpha)} \left(\ln \frac{\xi}{\tau}\right)^{\alpha i} \frac{h(\tau)}{\tau} d\tau \frac{d\xi}{\xi} = \\ &= \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + \alpha)} \int_a^t h(\tau) \int_{\tau}^t \left(\ln \frac{t}{\xi}\right)^{-\alpha} \left(\ln \frac{\xi}{\tau}\right)^{\alpha i + \alpha - 1} \frac{d\xi}{\xi} \frac{d\tau}{\tau} = \\ &= \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + \alpha)} \int_a^t h(\tau) \frac{\Gamma(1 - \alpha)\Gamma(\alpha i + \alpha)}{\Gamma(\alpha i + \alpha + 1 - \alpha)} \left(\ln \frac{t}{\tau}\right)^{\alpha i} \frac{d\tau}{\tau} \end{aligned}$$

Therefore,

$$\begin{aligned} D_a^{\alpha,\beta} h_0(t) &= t \frac{d}{dt} \left\{ \int_a^t h(\tau) \left[\sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + 1)} \left(\ln \frac{t}{\tau}\right)^{\alpha i} \right] \frac{d\tau}{\tau} \right\} = \\ &= t \frac{h(t)}{t} + \int_a^t h(\tau) \left[\sum_{i=1}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + 1)} \alpha i \left(\ln \frac{t}{\tau}\right)^{\alpha i - 1} \right] \frac{d\tau}{\tau} = \\ &= h(t) + \int_a^t h(\tau) \left[\sum_{j=0}^{\infty} \frac{\lambda^{j+1}}{\Gamma(\alpha j + \alpha)} \left(\ln \frac{t}{\tau}\right)^{\alpha j + \alpha - 1} \right] \frac{d\tau}{\tau} = \\ &= h(t) + \lambda \int_a^t h(\tau) \left(\ln \frac{t}{\tau}\right)^{\alpha - 1} E_{\alpha,\alpha} \left(\lambda \left(\ln \frac{t}{\tau}\right)^{\alpha} \right) \frac{d\tau}{\tau} d\xi = h(t) + \lambda h_0(t) \end{aligned}$$

Consequently,

$$D_a^{\alpha,\beta} h_0(t) = h(t) + \lambda h_0(t)$$

or the same

$$(D_a^{\alpha,\beta} - \lambda)h_0(t) = h(t).$$

In general, when $\alpha \in (m - 1, m]$ we can present function $D_a^{\alpha, \beta} h_0(t)$ in the following form:

$$D_a^{\alpha, \beta} h_0(t) = I_a^{\gamma_1} \delta^m I_a^\gamma (h_0)(t) = \frac{1}{\Gamma(\gamma_1)} \int_a^t \left(\ln \frac{t}{\tau}\right)^{\gamma_1 - 1} \delta g(\tau) \frac{d\tau}{\tau}, \tag{12}$$

where $g(\tau) = \delta^{m-1} I_a^\gamma [h_0](\tau)$.

Now we study the function $g(\tau)$. To do this, we transform the function $I_a^\gamma [h_0](\tau)$. By definition of the operator I_a^γ , we have

$$\begin{aligned} I_a^\alpha h_0(t) &= \frac{1}{\Gamma(\alpha)} \int_a^t \left(\ln \frac{t}{\tau}\right)^{\alpha - 1} h_0(\tau) \frac{d\tau}{\tau} = \\ &= \frac{1}{\Gamma(\alpha)} \int_a^t \left(\ln \frac{t}{\tau}\right)^{\alpha - 1} \int_a^\tau \left(\ln \frac{\tau}{\xi}\right)^{\alpha - 1} E_{\alpha, \alpha} \left(\lambda \left(\ln \frac{\tau}{\xi}\right)^\alpha\right) \frac{h(\xi)}{\xi} d\xi \frac{d\tau}{\tau} = \\ &= \sum_{i=0}^\infty \frac{\lambda^i}{\Gamma(\alpha i + \alpha)} \frac{1}{\Gamma(\alpha)} \int_a^t \left(\ln \frac{t}{\tau}\right)^{\alpha - 1} \int_a^\tau \left(\ln \frac{\tau}{\xi}\right)^{\alpha i + \alpha - 1} \frac{h(\xi)}{\xi} d\xi \frac{d\tau}{\tau} = \\ &= \sum_{i=0}^\infty \frac{\lambda^i}{\Gamma(\alpha i + \alpha)} \frac{1}{\Gamma(\alpha)} \int_a^t h(\xi) \int_\xi^t \left(\ln \frac{t}{\tau}\right)^{\alpha - 1} \left(\ln \frac{\tau}{\xi}\right)^{\alpha i + \alpha - 1} \frac{d\tau}{\tau} \frac{d\xi}{\xi} = \\ &= \int_a^t \left[\sum_{i=0}^\infty \frac{\lambda^i}{\Gamma(\alpha i + \alpha + \gamma)} \left(\ln \frac{t}{\xi}\right)^{\alpha i + \alpha - 1 + \gamma} \right] \frac{h(\xi)}{\xi} d\xi \end{aligned}$$

Further, since

$$\delta^{m-1} \left(\ln \frac{t}{\xi}\right)^{\alpha i + \alpha - 1 + \gamma} = (\alpha i + \alpha - 1 + \gamma) \dots (\alpha i + \alpha + \gamma - (m - 1))$$

and

$$\Gamma(\alpha i + \alpha + \gamma) = (\alpha i + \alpha + \gamma - 1) \dots (\alpha i + \alpha + \gamma - (m - 1)) \Gamma(\alpha i + \alpha + \gamma - (m - 1)),$$

then

$$D_a^{\alpha, \beta} h_0(t) = \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \frac{1}{\gamma_1} \left(\ln \frac{t}{\tau}\right)^{\gamma_1} \frac{d}{d\tau} g(\tau) d\tau \right\} =$$

$$= \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \left(\ln \frac{t}{\tau} \right)^{\gamma_1} g(\tau) d\tau \right\}.$$

Now it is obvious that $\lim_{\tau \rightarrow \infty} g(\tau) = 0$. Then by (12) as in the case $\alpha \in (0, 1)$ we get

$$\begin{aligned} D_a^{\alpha, \beta} h_0(t) &= \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \frac{1}{\gamma_1} \left(\ln \frac{t}{\tau} \right)^{\gamma_1} \frac{d}{d\tau} g(\tau) d\tau \right\} = \\ &= \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \left(\ln \frac{t}{\tau} \right)^{\gamma_1} g(\tau) d\tau \right\} \end{aligned}$$

Further,

$$\begin{aligned} &\int_a^t \left(\ln \frac{t}{\tau} \right)^{\gamma_1 - 1} g(\tau) d\tau = \\ &= \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(i\alpha + \alpha + \gamma - (m-1))} \int_a^t \left(\ln \frac{t}{\tau} \right)^{\gamma_1 - 1} \int_a^{\tau} \left(\ln \frac{\tau}{\xi} \right)^{\alpha i + \alpha + \gamma - m} \frac{h(\xi)}{\xi} d\xi \frac{d\tau}{\tau} = \\ &= \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(i\alpha + \alpha + \gamma - (m-1))} \int_a^t \frac{h(\xi)}{\xi} \int_{\xi}^t \left(\ln \frac{t}{\tau} \right)^{\gamma_1 - 1} \left(\ln \frac{\tau}{\xi} \right)^{\alpha i + \alpha + \gamma - m} \frac{d\tau}{\tau} d\xi = \\ &= \Gamma(\gamma_1) \sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + 1)} \int_a^t \left(\ln \frac{t}{\xi} \right)^{\alpha i} \frac{h(\xi)}{\xi} d\xi = \\ &= \Gamma(\gamma_1) \int_a^t \left[\sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + 1)} \left(\ln \frac{t}{\xi} \right)^{\alpha i} \right] \frac{h(\xi)}{\xi} d\xi \end{aligned}$$

Consequently,

$$\begin{aligned} D_a^{\alpha, \beta} h_0(t) &= t \frac{d}{dt} \left\{ \int_a^t \left[\sum_{i=0}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + 1)} \left(\ln \frac{t}{\xi} \right)^{\alpha i} \right] \frac{h(\xi)}{\xi} d\xi \right\} = \\ &= h(t) + \int_a^t \left[\sum_{i=1}^{\infty} \frac{\lambda^i}{\Gamma(\alpha i + 1)} \alpha i \left(\ln \frac{t}{\xi} \right)^{\alpha i - 1} \right] \frac{h(\xi)}{\xi} d\xi = \end{aligned}$$

$$= h(t) + \lambda \int_a^t \left[\sum_{j=0}^{\infty} \frac{\lambda^j}{\Gamma(\alpha j + \alpha)} \left(\ln \frac{t}{\xi} \right)^{\alpha j + \alpha - 1} \right] \frac{h(\xi)}{\xi} d\xi = h(t) + \lambda h_0(t).$$

So,

$$(D_a^{\alpha, \beta} - \lambda)h_0(t) = h(t).$$

Further, suppose $p = 0, 1, \dots$ $D_a^{\alpha, \beta} - \lambda$ and

$$E_{\alpha, \alpha}^p(\lambda, t) = \sum_{i=p}^{\infty} \lambda^{i-p} \binom{i}{p} \left(\ln \frac{t}{a} \right)^{\alpha i}.$$

It is obvious that

$$E_{\alpha, \alpha}^0(\lambda, t) = E_{\alpha, \alpha}(\lambda, t)$$

when $p = 0$.

Introduce the function:

$$h_p(t) = \int_a^t \left(\ln \frac{t}{\xi} \right)^{\alpha - 1} E_{\alpha, \alpha}^p \left(\lambda, \left(\ln \frac{t}{\xi} \right)^{\alpha} \right) h(\xi) \frac{d\xi}{\xi}.$$

Theorem 2. Let $m - 1 < \alpha \leq m, 0 \leq \beta \leq 1, s = 0, 1, \dots, m - 1$.

Then the functions $h_p(t), p = 0, 1, \dots$ form h -normed system concerning operator, i.e. the following equalities hold:

$$(D_a^{\alpha, \beta} - \lambda) h_0(t) = h(t), \tag{13}$$

$$(D_a^{\alpha, \beta} - \lambda) h_p(t) = h_{p-1}(t), \quad p \geq 1. \tag{14}$$

Proof. Since when $p = 0$ $E_{\alpha, \alpha}^0(\lambda, t) = E_{\alpha, \alpha}(\lambda, t)$ then we have proved equation (13) above. Let $p \geq 1$. As in the case $p = 0$ we present function $D_a^{\alpha, \beta} h_p(t)$ in the form:

$$D_a^{\alpha, \beta} h_p(t) = \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \frac{1}{\gamma_1} \left(\ln \frac{t}{\tau} \right)^{\gamma_1} \frac{d}{d\tau} f(\tau) d\tau \right\}, \tag{15}$$

where $\gamma_1 = \beta(m - \alpha), f(\tau) = \delta^{m-1} I_a^\gamma h_p(\tau)$. Further,

$$I_a^\alpha h_p(\tau) = \frac{1}{\Gamma(\gamma)} \int_a^\tau \left(\ln \frac{\tau}{\xi} \right)^{\gamma - 1} h_p(\xi) \frac{d\xi}{\xi} =$$

$$\begin{aligned}
 &= \frac{1}{\Gamma(\gamma)} \int_a^\tau \left(\ln \frac{\tau}{\xi}\right)^{\gamma-1} \int_a^\xi \left[\sum_{i=p}^\infty \frac{\lambda^{i-p}}{\Gamma(\alpha i + \alpha)} \binom{i}{p} \left(\ln \frac{\xi}{z}\right)^{\alpha i + \alpha - 1} \right] \frac{h(z) dz}{z \xi} = \\
 &= \sum_{i=p}^\infty \frac{\lambda^{i-p}}{\Gamma(\alpha i + \alpha)} \binom{i}{p} \frac{1}{\Gamma(\gamma)} \int_a^\tau h(z) \int_z^\tau \left(\ln \frac{\tau}{\xi}\right)^{\gamma-1} \left(\ln \frac{\xi}{z}\right)^{\alpha i + \alpha - 1} \frac{d\xi dz}{\xi z} = \\
 &= \int_a^\tau \left[\sum_{i=p}^\infty \frac{\lambda^{i-p}}{\Gamma(\alpha i + \alpha + \gamma)} \binom{i}{p} \left(\ln \frac{\tau}{z}\right)^{\alpha i + \alpha - 1 + \gamma} \right] h(z) \frac{dz}{z}
 \end{aligned}$$

Hence for a function $f(\tau)$ we obtain

$$f(\tau) = \delta^m I_a^\gamma [h_p](\tau) = \int_a^\tau \left[\sum_{i=p}^\infty \frac{\lambda^{i-p}}{\Gamma(\alpha i + \alpha + \gamma - (m-1))} \left(\ln \frac{\tau}{z}\right)^{\alpha i + \alpha + \gamma - m} \right] h(z) \frac{dz}{z}.$$

Consequently, integrating by parts in (15), we get

$$D_a^{\alpha, \beta} h_p(t) = \frac{1}{\Gamma(\gamma_1)} t \frac{d}{dt} \left\{ \int_a^t \left(\ln \frac{t}{\tau}\right)^{\gamma_1 - 1} f(\tau) \frac{d\tau}{\tau} \right\}.$$

Further, as in the case $p = 0$ we have

$$\int_a^t \left(\ln \frac{t}{\tau}\right)^{\gamma_1 - 1} f(\tau) \frac{d\tau}{\tau} = \Gamma(\gamma_1) \int_a^t \left[\sum_{i=p}^\infty \frac{\lambda^{i-p}}{\Gamma(\alpha i + 1)} \binom{i}{p} \left(\ln \frac{t}{\xi}\right)^{\alpha i} \right] \frac{h(\xi)}{\xi} d\xi.$$

It means that

$$\begin{aligned}
 D_a^{\alpha, \beta} h_p(t) &= \int_a^t \left[\sum_{i=p}^\infty \frac{\lambda^{i-p}}{\Gamma(\alpha i)} \binom{i}{p} \left(\ln \frac{t}{\xi}\right)^{\alpha i - 1} \right] \frac{h(\xi)}{\xi} d\xi = \\
 &= \int_a^t \left[\sum_{j=p-1}^\infty \frac{\lambda^{j-(p-1)}}{\Gamma(\alpha j + \alpha)} \binom{j+1}{p} \left(\ln \frac{t}{\xi}\right)^{\alpha j + \alpha - 1} \right] \frac{h(\xi)}{\xi} d\xi.
 \end{aligned}$$

Then

$$\begin{aligned}
 &D_a^{\alpha, \beta} h_p(t) - h_{p-1}(t) = \\
 &= \int_a^t \left\{ \sum_{i=p-1}^\infty \left[\binom{i+1}{p} - \binom{i}{p-1} \right] \frac{\lambda^{i-(p-1)}}{\Gamma(\alpha i + \alpha)} \left(\ln \frac{t}{\xi}\right)^{\alpha i + \alpha - 1} \right\} \frac{h(\xi)}{\xi} d\xi =
 \end{aligned}$$

$$= \int_a^t \left\{ \sum_{i=p}^{\infty} \binom{i}{p} \frac{\lambda^{i-(p-1)}}{\Gamma(\alpha i + \alpha)} \left(\ln \frac{t}{\xi} \right)^{\alpha i + \alpha - 1} \right\} \frac{h(\xi)}{\xi} d\xi = \lambda h_p(t).$$

Therefore,

$$D_a^{\alpha, \beta} h_p(t) - h_{p-1}(t) = \lambda h_p(t),$$

i.e. for all $p = 1, 2, \dots$ the following equality holds:

$$\left(D_a^{\alpha, \beta} - \lambda \right) h_p(t) = h_{p-1}(t).$$

Theorem is proved. Theorems 1 and 2 imply the following basic statement:

Theorem 3. *Let $m - 1 < \alpha \leq m, m = 1, 2, \dots, 0 \leq \beta \leq 1$. Then the function*

$$y(t) = \sum_{s=0}^{m-1} \sum_{p=0}^{n-1} C_{s,p} \Phi_{\alpha, \beta, s}^p(\lambda, t) + \int_a^t \left(\ln \frac{t}{\xi} \right)^{\alpha-1} E_{\alpha, \alpha}^{n-1} \left(\lambda, \left(\ln \frac{t}{\xi} \right)^\alpha \right) h(\xi) \frac{d\xi}{\xi}$$

is a solution of equation (1), where $C_{s,p}$ are constants.

Acknowledgments

This paper is financial supported by the grant of MES RK No.0713/GF.

References

- [1] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, North-Holland (2006).
- [2] M.D. Qassim, K.M. Furati, N.-E. Tatar, On a differential equation involving hilfer-Hadamard fractional derivative, *Abstract and Applied Analysis* (2012), Article ID 391062, 17.
- [3] R. Hilfer, Y. Luchko, Z. Tomovski, Operational method the solution of fractional differential equations with generalized Rieman-Lioville fractional derivatives, *Fractional Calculus and Applied Analysis*, **12**, No. 3 (2009), 299-318.
- [4] B.A. Bondarenko, *Operator Algorithms in Differential Equations*, Tashkent, "FAN" (1984), In Russian.

- [5] K.M. Shinaliyev, B.Kh. Turmetov, S.A. Umarov, Fractional operator algorithm method for construction of solutions of fractional order differential equations, *Fractional Calculus and Applied Analysis*, **15**, No. 2 (2012), 267-281.
- [6] B.Kh. Turmetov, K.M. Shinaliyev, On an operator method of construction the solution of differential equations of the fractional order, *Vestnik of L. Gumilyev Eurasian National University. Series for Natural and Engineering Sciences*, **81**, No. 2 (2011), 10-17.